

PCT / SE 2004 / 001941

20-12-2004

PA 1246077

REC'D 18 JAN 2005

WIPO

PCT

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

November 08, 2004

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.


APPLICATION NUMBER: 60/531,710

FILING DATE: December 22, 2003

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS



H. L. Jackson
H. L. JACKSON

Certifying Officer

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Express Mail Label No. ET971611053US

INVENTOR(S)

Given Name (first and middle [if any]) Edon	Family Name or Surname PHILLIPS	Residence (City and either State or Foreign Country) Wilmington, DE
--	------------------------------------	---

Additional inventors are being named on the _____ separately numbered sheets attached hereto

TITLE OF THE INVENTION (500 characters max)

NICOTINIC ACETYLCHOLINE RECEPTOR LIGANDS

Direct all correspondence to:

CORRESPONDENCE ADDRESS
☒ Customer Number: 22466
 OR

<input type="checkbox"/> Firm or Individual Name					
Address					
Address					
City		State		ZIP	
Country		Telephone		Fax	

ENCLOSED APPLICATION PARTS (check all that apply)

<input checked="" type="checkbox"/> Specification Number of Pages 22	<input type="checkbox"/> CD(s), Number _____
<input type="checkbox"/> Drawing(s) Number of Sheets _____	<input type="checkbox"/> Other (specify) _____
<input type="checkbox"/> Application Data Sheet. See 37 CFR 1.76	

METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT

<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27.	FILING FEE Amount (\$) 160.00
<input type="checkbox"/> A check or money order is enclosed to cover the filing fees.	
<input checked="" type="checkbox"/> The Director is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number: 26-0166	
<input type="checkbox"/> Payment by credit card. Form PTO-2038 is attached.	

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

<input checked="" type="checkbox"/> No.
<input type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are: _____

[Page 1 of 1]

Respectfully submitted,

SIGNATURE

K F Mitchell

TYPED or PRINTED NAME Kenneth F. Mitchell

TELEPHONE 302 886 7466

Date December 22, 2003

REGISTRATION NO. 42,007

(If appropriate)

Docket Number: 101332-1 US

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

This collection of information is required by 37 CFR 1.51. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Mail Stop Provisional Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

NICOTINIC ACETYLCHOLINE RECEPTOR LIGANDS

FIELD OF THE INVENTION

This invention relates to diazabicyclo-octyl amides or pharmaceutically-acceptable salts thereof, processes for preparing them, pharmaceutical compositions containing them and their use in therapy. The invention also relates to compounds that are ligands for nicotinic acetylcholine receptors (nAChRs).


BACKGROUND OF THE INVENTION

The use of compounds which bind nicotinic acetylcholine receptors in the treatment of a range of disorders involving reduced cholinergic function such as Alzheimer's disease, cognitive or attention disorders, anxiety, depression, smoking cessation, neuroprotection, schizophrenia, analgesia, Tourette's syndrome, and Parkinson's disease has been discussed in McDonald et al. (1995) "Nicotinic Acetylcholine Receptors: Molecular Biology, Chemistry and Pharmacology", Chapter 5 in Annual Reports in Medicinal Chemistry, vol. 30, pp. 41-50, Academic Press Inc., San Diego, CA; and in Williams et al. (1994) "Neuronal Nicotinic Acetylcholine Receptors," Drug News & Perspectives, vol. 7, pp. 205-223.

SUMMARY OF THE INVENTION

This invention concerns nicotinic acetylcholine receptor-active compounds of formula

I:



wherein:

D is selected from oxygen, sulfur or $N(R^1)_2$;

Ar^1 is selected from a 5- or 6-membered aromatic or heteroaromatic ring having 0, 1 or 2 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms, or selected from an 8-, 9- or 10-membered fused aromatic or heteroaromatic ring system having 0, 1, 2 or 3 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms;

E is a single bond, -O, -S, or $-NR^2$;

G is selected from hydrogen, C₁-C₄alkoxy or Ar², where Ar² is a 5- or 6-membered aromatic or heteroaromatic ring having 0, 1 or 2 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms;

where each Ar¹ or Ar² moiety independently is unsubstituted or has 1, 2 or 3 substituents selected from -R³, -C₁-C₆alkyl, -C₂-C₆alkenyl, -C₂-C₆alkynyl, halogen, -CN, -NO₂, -CF₃, -S(O)_nR³, -NR²R³, -CH₂NR²R³, -OR³, -CH₂OR³ or -CO₂R⁴;

R¹, R² and R³ are independently selected at each occurrence from hydrogen, -C₁-C₄alkyl, aryl, heteroaryl, -C(O)R⁴, -C(O)NHR⁴, -CO₂R⁴ or -SO₂R⁴, or

R² and R³ in combination is -(CH₂)_jG(CH₂)_k- wherein G is oxygen, sulfur, NR⁴, or a bond;

j is 2, 3 or 4;

k is 0, 1 or 2;


n is 0, 1 or 2, and

R⁴ is independently selected at each occurrence from hydrogen, -C₁-C₄alkyl, aryl, or heteroaryl.

The invention also encompasses stereoisomers, enantiomers, *in vivo*-hydrolysable precursors and pharmaceutically-acceptable salts of compounds of formula I, pharmaceutical compositions and formulations containing them, methods of using them to treat diseases and conditions either alone or in combination with other therapeutically-active compounds or substances, processes and intermediates used to prepare them, uses of them as medicaments, uses of them in the manufacture of medicaments and uses of them for diagnostic and analytic purposes.

DETAILED DESCRIPTION OF THE INVENTION

Compound of the invention are those according to formula I:



wherein:

D is selected from oxygen, sulfur or N(R¹)₂;

Ar¹ is selected from a 5- or 6-membered aromatic or heteroaromatic ring having 0, 1 or 2 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms, or selected from an 8-, 9- or 10-membered fused aromatic or heteroaromatic ring system having 0, 1, 2 or 3 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms;

5 E is a single bond, -O, -S, or -NR²;

G is selected from hydrogen, C₁-C₄alkoxy or Ar², where Ar² is a 5- or 6-membered aromatic or heteroaromatic ring having 0, 1 or 2 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms;

10 where each Ar¹ or Ar² moiety independently is unsubstituted or has 1, 2 or 3 substituents selected from -R³, -C₁-C₆alkyl, -C₂-C₆alkenyl, -C₂-C₆alkynyl, halogen, -CN, -NO₂, -CF₃, -S(O)_nR³, -NR²R³, -CH₂NR²R³, -OR³, -CH₂OR³ or -CO₂R⁴;

R¹, R² and R³ are independently selected at each occurrence from hydrogen, -C₁-C₄alkyl, aryl, heteroaryl, -C(O)R⁴, -C(O)NHR⁴, -CO₂R⁴ or -SO₂R⁴, or

15 R² and R³ in combination is -(CH₂)_jG(CH₂)_k- wherein G is oxygen, sulfur, NR⁴, or a bond;

j is 2, 3 or 4;

k is 0, 1 or 2;

n is 0, 1 or 2, and

20 R⁴ is independently selected at each occurrence from hydrogen, -C₁-C₄alkyl, aryl, or heteroaryl,

and stereoisomers, enantiomers, *in vivo*-hydrolysable precursors and pharmaceutically-acceptable salts thereof.

More particular compounds are those of formula I wherein:

D is oxygen;

25 Ar¹ is selected from phenyl or a 5-membered heteroaromatic ring having 0 or 1 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms, or selected from a 9-membered fused aromatic or heteroaromatic ring system having 0, 1, 2 or 3 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms;;

wherein:

30 E is a single bond;

G is selected from hydrogen, methoxy or Ar², where Ar² is selected from a 6-membered aromatic or heteroaromatic ring having 0 or 1 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms;

where each Ar¹ or Ar² moiety independently is unsubstituted or has 1, 2 or 3 substituents selected from halogen, -CN, -NO₂, -CF₃, -CH₃ or -C₂H₅; and stereoisomers, enantiomers, *in vivo*-hydrolysable precursors and pharmaceutically-acceptable salts thereof.

Still more particular compounds are those of formula I wherein:

D is oxygen;

Ar¹ is selected from phenyl, furanyl, thiophenyl or 1-methyl-1H-pyrrolyl;

E is a single bond;

G is selected from hydrogen, methoxy, phenyl or pyridyl, and

Ar¹ bears 1 halogen substituent;

and stereoisomers, enantiomers, *in vivo*-hydrolysable precursors and pharmaceutically-acceptable salts thereof.

Other particular compounds of the invention include those of formula I wherein E represents a single bond; or an enantiomer thereof, and pharmaceutically-acceptable salts thereof.

Particular compounds of the invention are those of formula I wherein Ar¹ is furanyl or thiophenyl having optional substituents as defined herein.

Particular compounds of the invention are those described herein and pharmaceutically-acceptable salts thereof.

In a further aspect the invention encompasses compounds according to formula I wherein one or more of the atoms is a radioisotope of the same element. In a particular form of this aspect of the invention the compound of formula I is labeled with tritium. Such radio-labeled compounds are synthesized either by incorporating radio-labeled starting materials or, in the case of tritium, exchange of hydrogen for tritium by known methods. Known methods include (1) electrophilic halogenation, followed by reduction of the halogen in the presence of a tritium source, for example, by hydrogenation with tritium gas in the presence of a palladium catalyst, or (2) exchange of hydrogen for tritium performed in the presence of tritium gas and a suitable organometallic (e.g. palladium) catalyst.

Compounds of the invention labeled with tritium are useful for the discovery of novel medicinal compounds which bind to and modulate the activity, by agonism, partial agonism, or antagonism, of the $\alpha 7$ nicotinic acetylcholine receptor. Such tritium-labeled compounds may be used in assays that measure the displacement of a such compounds to assess the binding of ligand that bind to $\alpha 7$ nicotinic acetylcholine receptors.

In another aspect the invention relates to compounds according to formula I and their use in therapy and to compositions containing them.

In another aspect the invention encompasses the use of compounds according to formula I for the therapy of diseases mediated through the action of nicotinic acetylcholine receptors. A more particular aspect of the invention relates to the use of compounds of formula I for the therapy of diseases mediated through the action of $\alpha 7$ nicotinic acetylcholine receptors.

Another aspect of the invention encompasses a method of treatment or prophylaxis of diseases or conditions in which activation of the $\alpha 7$ nicotinic receptor is beneficial which method comprises administering a therapeutically-effective amount of a compound of the invention to a subject suffering from said disease or condition.

One embodiment of this aspect of the invention is a method of treatment or prophylaxis, wherein the disorder is anxiety, schizophrenia, mania or manic depression.

Another embodiment of this aspect of the invention is a method of treatment or prophylaxis of neurological disorders, psychotic disorders or intellectual impairment disorders, which comprises administering a therapeutically effective amount of a compound of the invention.

Another embodiment of this aspect of the invention is a method of treatment or prophylaxis, wherein the disorder is Alzheimer's disease, learning deficit, cognition deficit, attention deficit, memory loss, or Attention Deficit Hyperactivity Disorder.

Another embodiment of this aspect of the invention is a method of treatment or prophylaxis, wherein the disorder is Parkinson's disease, Huntington's disease, Tourette's syndrome, or neurodegenerative disorders in which there is loss of cholinergic synapses.

Another embodiment of this aspect of the invention is a method of treatment or prophylaxis of jetlag, nicotine addiction, craving, pain, and for ulcerative colitis, which comprises administering a therapeutically effective amount of a compound of the invention.

Yet another embodiment of this aspect of the invention is a method for inducing the cessation of smoking which comprises administering an effective amount of a compound of the invention.

Another embodiment of this aspect of the invention is a pharmaceutical composition
5 comprising a compound of the invention and a pharmaceutically-acceptable diluent, lubricant or carrier.

A further aspect of the invention relates to a pharmaceutical composition useful for treating or preventing a condition or disorder mentioned herein arising from dysfunction of nicotinic acetylcholine receptor neurotransmission in a mammal, preferably a human,
10 comprising an amount of a compound of formula I, an enantiomer thereof or a pharmaceutically-acceptable salt thereof, effective in treating or preventing such disorder or condition, and pharmaceutically-acceptable additives carrier.

Another embodiment of this aspect of the invention relates to use of a pharmaceutical composition of the invention for the treatment, amelioration or prophylaxis of human diseases
15 or conditions in which activation of the $\alpha 7$ nicotinic receptor is beneficial.

Another embodiment of this aspect of the invention is the use of the pharmaceutical composition of the invention for the treatment or prophylaxis of neurological disorders, psychotic disorders or intellectual impairment disorders.

Another embodiment of this aspect of the invention is the use of the pharmaceutical
20 composition of the invention for the treatment or prophylaxis of Alzheimer's disease, learning deficit, cognition deficit, attention deficit, memory loss, Attention Deficit Hyperactivity Disorder, anxiety, schizophrenia, or mania or manic depression, Parkinson's disease, Huntington's disease, Tourette's syndrome, neurodegenerative disorders in which there is loss of cholinergic synapse, jetlag, cessation of smoking, nicotine addiction including
25 that resulting from exposure to products containing nicotine, craving, pain, and for ulcerative colitis.

A further aspect of the invention is the use of a compound according to the invention, an enantiomer thereof or a pharmaceutically-acceptable salt thereof, in the manufacture of a medicament for the treatment or prophylaxis of the diseases or conditions mentioned herein.

30 Another embodiment of this aspect of the invention is the use of a compound of the invention in the manufacture of a medicament for the treatment or prophylaxis of human diseases or conditions in which activation of the $\alpha 7$ nicotinic receptor is beneficial.

Another embodiment of this aspect of the invention is the use of a compound of the invention in the manufacture of a medicament for the treatment or prophylaxis of neurological disorders, psychotic disorders or intellectual impairment disorders.

5 Another embodiment of this aspect of the invention is the use of a compound of the invention in the manufacture of a medicament for treatment or prophylaxis of Alzheimer's disease, learning deficit, cognition deficit, attention deficit, memory loss or Attention Deficit Hyperactivity Disorder.

10 Another embodiment of this aspect of the invention is the use of a compound of the invention in the manufacture of a medicament for treatment or prophylaxis of anxiety, schizophrenia, or mania or manic depression.

Another embodiment of this aspect of the invention is the use of a compound of the invention in the manufacture of a medicament for treatment or prophylaxis of Parkinson's disease, Huntington's disease, Tourette's syndrome, or neurodegenerative disorders in which there is loss of cholinergic synapses.

15 Another embodiment of this aspect of the invention is the use of a compound as described above in the manufacture of a medicament for the treatment or prophylaxis of jetlag, pain, or ulcerative colitis.

20 Another aspect of the invention relates to the use of a compound of the invention in the manufacture of a medicament for facilitating the cessation of smoking or the treatment of nicotine addiction or craving including that resulting from exposure to products containing nicotine.

25 For the uses, methods, medicaments and pharmaceutical compositions mentioned herein the amount of compound used and the dosage administered will, of course, vary with the compound employed, the mode of administration and the treatment desired. However, in general, satisfactory results are obtained when the compounds of the invention are administered at a daily dosage of from about 0.1 mg to about 20 mg/kg of animal body weight. Such doses may be given in divided doses 1 to 4 times a day or in sustained release form. For man, the total daily dose is in the range of from 5 mg to 1,400 mg, more preferably from 10 mg to 100 mg, and unit dosage forms suitable for oral administration comprise from 30 2 mg to 1,400 mg of the compound admixed with a solid or liquid pharmaceutical carriers, lubricants and diluents.

The compounds of formula I, an enantiomer thereof, and pharmaceutically-acceptable salts thereof, may be used on their own or in the form of appropriate medicinal preparations for enteral or parenteral administration. According to a further aspect of the invention, there is provided a pharmaceutical composition including preferably less than 80% and more
5 preferably less than 50% by weight of a compound of the invention in admixture with an inert pharmaceutically-acceptable diluent, lubricant or carrier.

Examples of diluents, lubricants and carriers are:

- for tablets and dragees: lactose, starch, talc, stearic acid;
- for capsules: tartaric acid or lactose;
- 10 – for injectable solutions: water, alcohols, glycerin, vegetable oils;
- for suppositories: natural or hardened oils or waxes.

There is also provided a process for the preparation of such a pharmaceutical composition which process comprises mixing the ingredients.

Compounds according to the invention are agonists of nicotinic acetylcholine
15 receptors. While not being limited by theory, it is believed that agonists of the $\alpha 7$ nicotinic acetylcholine receptor (nAChR) subtype are useful in the treatment or prophylaxis of neurological disorders, psychotic disorders and intellectual impairment disorders, and to have advantages over compounds which are or are also agonists of the $\alpha 4$ nAChR subtype. Therefore, compounds which are selective for the $\alpha 7$ nAChR subtype are preferred. The
20 compounds of the invention are indicated as pharmaceuticals, in particular in the treatment or prophylaxis of neurological disorders, psychotic disorders and intellectual impairment disorders. Examples of psychotic disorders include schizophrenia, mania and manic depression, and anxiety. Examples of intellectual impairment disorders include Alzheimer's disease, learning deficit, cognition deficit, attention deficit, memory loss, and Attention
25 Deficit Hyperactivity Disorder. The compounds of the invention may also be useful as analgesics in the treatment of pain, chronic pain, and in the treatment or prophylaxis of Parkinson's disease, Huntington's disease, Tourette's syndrome, and neurodegenerative disorders in which there is loss of cholinergic synapses.

Compounds of the invention may further useful for the treatment or prophylaxis of
30 jetlag, for use in inducing the cessation of smoking, craving, and for the treatment or prophylaxis of nicotine addiction including that resulting from exposure to products containing nicotine.

It is also believed that compounds according to the invention are useful in the treatment and prophylaxis of ulcerative colitis.

The compounds of the invention have the advantage that they may be less toxic, be more efficacious, be longer acting, have a broader range of activity, be more potent, produce
5 fewer side effects, are more easily absorbed or have other useful pharmacological properties.

The compounds of formula I exist in tautomeric or enantiomeric forms, all of which are included within the scope of the invention. The various optical isomers may be isolated by separation of a racemic mixture of the compounds using conventional techniques, e.g. fractional crystallization, or chiral HPLC. Alternatively the individual enantiomers may be
10 made by reaction of the appropriate optically active starting materials under reaction conditions which will not cause racemization.

As used herein, unless otherwise indicated, "C₁₋₄alkyl" includes but is not limited to methyl, ethyl, *n*-propyl, *n*-butyl, *i*-propyl, *i*-butyl, *t*-butyl, *s*-butyl moieties, whether alone or part of another group, C₁₋₄alkyl groups may be straight-chained or branched, and C₃₋₄ alkyl
15 groups include the cyclic alkyl moieties cyclopropyl and cyclobutyl.

As used herein, unless otherwise indicated, "C₂₋₄alkenyl" includes but is not limited to 1-propenyl, 2-propenyl, 1-butenyl, 2-butenyl and 3-butenyl.

As used herein, unless otherwise indicated, "C₂₋₄alkynyl" includes but is not limited to ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butyne and 3-butyne.

20 As used herein, unless otherwise indicated, aryl refers to a phenyl ring which may have 1, 2 or 3 substituents selected from: halogen, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, C₁₋₄alkyl, CN, NO₂, and CF₃.

As used herein, unless otherwise indicated, heteroaryl refers to a 5- or 6-membered aromatic or heteroaromatic ring having 1, 2 or 3 heteroatoms selected from nitrogen oxygen
25 and sulfur, provided that heteroaromatic rings contains at least one nitrogen, oxygen, or sulfur atom.

As used herein, unless otherwise indicated, halogen refers to fluorine, chlorine, bromine, or iodine.

Where necessary, hydroxy, amino, or other reactive groups may be protected using a
30 protecting group as described in the standard text "Protecting groups in Organic Synthesis", 3rd Edition (1999) by Greene and Wuts.

Unless otherwise stated, reactions are conducted under an inert atmosphere, preferably under a nitrogen atmosphere and are usually conducted at a pressure of about one to about three atmospheres, preferably at ambient pressure (about one atmosphere).

The compounds of the invention and intermediates may be isolated from their reaction mixtures by standard techniques.

Acid addition salts of the compounds of formula I which may be mentioned include salts of mineral acids, for example the hydrochloride and hydrobromide salts; and salts formed with organic acids such as formate, acetate, maleate, benzoate, tartrate, and fumarate salts.

Acid addition salts of compounds of formula I may be formed by reacting the free base or a salt, enantiomer or protected derivative thereof, with one or more equivalents of the appropriate acid. The reaction may be carried out in a solvent or medium in which the salt is insoluble or in a solvent in which the salt is soluble, e.g., water, dioxane, ethanol, tetrahydrofuran or diethyl ether, or a mixture of solvents, which may be removed in vacuum or by freeze drying. The reaction may be a metathetical process or it may be carried out on an ion exchange resin.

The compounds of formula I exist in tautomeric or enantiomeric forms, all of which are included within the scope of the invention. The various optical isomers may be isolated by separation of a racemic mixture of the compounds using conventional techniques, e.g. fractional crystallisation, or chiral HPLC. Alternatively the individual enantiomers may be made by reaction of the appropriate optically active starting materials under reaction conditions which will not cause racemisation.

Pharmacology

The pharmacological activity of the compounds of the invention may be measured in the tests set out below:

Test A - Assay for affinity at α_7 nAChR subtype

¹²⁵I- α -Bungarotoxin (BTX) binding to rat hippocampal membranes.

Rat hippocampi were homogenized in 20 volumes of cold homogenisation buffer (HB: concentrations of constituents (mM): tris(hydroxymethyl)aminomethane 50; MgCl₂ 1; NaCl 120; KCl 5; pH 7.4). The homogenate was centrifuged for 5 minutes at 1000 xg, the supernatant was saved and the pellet re-extracted. The pooled supernatants were centrifuged for 20 minutes at 12000 xg, washed, and re-suspended in HB. Membranes (30–80 μ g) were

incubated with 5 nM [125 I] α -BTX, 1 mg/mL BSA (bovine serum albumin), test drug, and either 2 mM CaCl₂ or 0.5 mM EGTA [ethylene glycol-bis(β -aminoethylether)] for 2 hours at 21 °C, and then filtered and washed 4 times over Whatman glass fibre filters (thickness C) using a Brandel cell harvester. Pre-treating the filters for 3 hours with 1% (BSA/0.01% PEI (polyethyleneimine) in water was critical for low filter blanks (0.07% of total counts per minute). Non-specific binding was described by 100 μ M (-)-nicotine, and specific binding was typically 75%.

Test B - Assay for affinity to the α_4 nAChR subtype

[3 H]-(-)-nicotine binding.

Using a procedure modified from Martino-Barrows and Kellar (Mol Pharm (1987) 31:169-174), rat brain (cortex and hippocampus) was homogenised as in the [125 I] α -BTX binding assay, centrifuged for 20 minutes at 12,000 xg, washed twice, and then re-suspended in HB containing 100 μ M diisopropyl fluorophosphate. After 20 minutes at 4 °C, membranes (approximately 0.5 mg) were incubated with 3 nM [3 H]-(-)-nicotine, test drug, 1 μ M atropine, and either 2 mM CaCl₂ or 0.5 mM EGTA for 1 hour at 4 °C, and then filtered over Whatman glass fibre filters (thickness C) (pre-treated for 1 hour with 0.5% PEI) using a Brandel cell harvester. Non-specific binding was described by 100 μ M carbachol, and specific binding was typically 84%.

Binding data analysis for Tests A and B

IC₅₀ values and pseudo Hill coefficients (n_H) were calculated using the non-linear curve fitting program ALLFIT (DeLean A, Munson P J and Rodbard D (1977) Am. J. Physiol., 235:E97-E102). Saturation curves were fitted to a one site model, using the non-linear regression program ENZFITTER (Leatherbarrow, R.J. (1987)), yielding K_D values of 1.67 and 1.70 nM for the 125 I- α -BTX and [3 H]-(-)-nicotine ligands respectively. K_i values were estimated using the general Cheng-Prusoff equation:

$$K_i = [IC_{50}] / ((2 + ([\text{ligand}] / K_D)^{n/n-1})$$


where a value of $n=1$ was used whenever $n_H < 1.5$ and a value of $n=2$ was used when $n_H \geq 1.5$. Samples were assayed in triplicate and were typically $\pm 5\%$. K_i values were determined using 6 or more drug concentrations. The compounds of the invention are

compounds with binding affinities (K_i) of less than 10 μ M in either Test A or Test B, indicating that they are expected to have useful therapeutic activity.

The compounds of the invention have the advantage that they may be less toxic, be more efficacious, be longer acting, have a broader range of activity, be more potent, produce fewer side effects, are more easily absorbed or have other useful pharmacological properties.


Intermediate 1: 1,4-Diazabicyclo[3.2.1]octane

a) 3-Oxo-piperazin-2-yl-acetic acid ethyl ester



3-Oxo-piperazin-2-yl-acetic acid ethyl ester was prepared according to the procedure described by S. Gubert, et. al. (*J. Het. Chem.*, **30**, 1993, 275-276).


b) 2-Piperazin-2-yl-ethanol



To a mixture of 3-oxo-piperazin-2-yl-acetic acid ethyl ester (2.0 g, 10.74 mmol) in 50 mL of dry THF cooled in an ice bath, was added LAH (1M solution in THF, 20.0 mL, 20.0 mmol) dropwise with stirring under N_2 . When addition was complete (c. 10 min), the reaction mixture was refluxed for 3½ h, then cooled in an ice bath. Water (5 mL) was cautiously added with stirring. After stirring for ½ h, the mixture was filtered through a fritted funnel and the collected salts were washed with hot EtOH. The filtrates were combined, dried over $MgSO_4$, filtered and solvents removed *in vacuo*. The residue was treated with hot $CHCl_3$, filtered and the $CHCl_3$ was evaporated to give a pale yellow oil. The product was obtained in quantitative yield and carried forward without further purification.


1H NMR (300.132 MHz, $CDCl_3$) δ 3.82 - 3.78 (m, 1H), 2.98 - 2.63 (m, 5H), 2.45 - 2.36 (m, 1H), 1.62 - 1.53 (m, 3H), 1.66 (bs, 2H), 1.13 (bs, 1H).

c) 1,4-Diazabicyclo[3.2.1]octane dihydrochloride salt




The title compound, 1,4-diazabicyclo[3.2.1]octane was prepared as a dihydrochloride salt from 2-piperazin-2-yl-ethanol according to the procedure described by P. A. Sturn et. al. (J. Med. Chem., 20 (10), 1977, 1333-1337.

Example 1: (1,4-Diazabicyclo[3.2.1]oct-4-yl)-(5-pyridin-3-yl-thiophen-2-yl)-methanone



To a stirred solution of 5-(2-pyridyl)thiophene-2-carboxylic acid (45.0 mg, 0.22 mmol), O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate TBTU (71.0 mg, 0.22 mmol), and 1-hydroxybenzotriazole hydrate (30.0 mg, 0.22 mmol) in DMF (2 mL), was added diisopropylethylamine (0.05 mL, 0.29 mmol). After 5 min, a mixture of 1,4-diazabicyclo[3.2.1]octane dihydrochloride salt (40.0 mg, 0.22 mmol) and 0.1 mL DIEA (0.1 mL, 0.59 mmol) in DMF (1 mL) was added. The reaction mixture was stirred at room temperature overnight. The reaction mixture was then partitioned between EtOAc and 5% Na₂CO₃. The layers were separated and the aqueous phase was extracted with EtOAc. The organic extracts were combined, dried over MgSO₄, filtered and concentrated *in vacuo*. The residue was chromatographed on silica gel using a gradient of 100:0 to 95:5 CHCl₃:MeOH. The product was obtained as an off-white solid (39 mg, 60 %). MS (APCI+) 300 [M+1]⁺. ¹H NMR (300.132 MHz, CDCl₃) δ 8.89 (s, 1H), 8.58 (d, *J* = 4.2 Hz, 1H), 7.87 (dt, *J* = 8.0 Hz, *J* = 1.9 Hz, 1H), 7.34 (dd, *J* = 3.1 Hz, *J* = 4.9 Hz, 1H), 7.30 (q, *J* = 7.9 Hz, 2H), 5.04 (s, 1H), 4.11 (dd, *J* = 13.9 Hz, *J* = 5.2 Hz, 1H), 3.43 (t, *J* = 10.7 Hz, 1H), 3.23 - 3.04 (m, 2H), 2.88 (dd, *J* = 4.7 Hz, *J* = 13.8 Hz, 1H), 2.77 (d, *J* = 11.0 Hz, 1H), 2.55-2.34 (m, 2H), 2.19 - 1.97 (m, 2H).


Example 2: (1,4-Diazabicyclo[3.2.1]oct-4-yl)-(5-phenyl-thiophen-2-yl)-methanone



By the process described in Example 1, diazabicyclo[3.2.1]octane dihydrochloride salt with 5-phenyl-thiophene-2-carboxylic acid to afford the title compound as an amber gum. MS (APCI+) 299 [M+1]⁺. ¹H NMR (300.132 MHz, CDCl₃) δ 7.61 (dt, *J* = 7.5, 1.7 Hz, 2H),


7.40 (tt, $J = 7.3, 1.6$ Hz, 2H), 7.34 (dt, $J = 7.2, 1.5$ Hz, 1H), 7.28 (d, $J = 3.6$ Hz, 1H), 7.23 (d, $J = 3.6$ Hz, 1H), 4.98 (m, 1H), 4.03 (dd, $J = 13.5, 4.8$ Hz, 1H), 3.37 (m, 1H), 3.09 (d, $J = 13.1$ Hz, 1H), 3.05 (t, $J = 8.0$ Hz, 3H), 2.77 (dd, $J = 13.4, 4.2$ Hz, 1H), 2.63 (d, $J = 11.6$ Hz, 1H), 2.06 - 1.97 (m, 2H).

5 **Example 3:** [5-(4-Chloro-phenyl)-furan-2-yl]-(1,4-diazabicyclo[3.2.1]oct-4-yl)-methanone




By the process described in Example 1, diazabicyclo[3.2.1]octane dihydrochloride salt was reacted with 5-(4-chloro-phenyl)-furan-2-carboxylic acid to afford the title compound as a gum. MS (APCI+) 317/319 $[M+1]^+$. ^1H NMR (300.132 MHz, CDCl_3) δ 7.61 (d, $J = 8.6$ Hz, 2H), 7.38 (d, $J = 8.6$ Hz, 2H), 7.06 (d, $J = 3.5$ Hz, 1H), 6.71 (d, $J = 3.5$ Hz, 1H), 5.08 (m, 1H), 4.13 (dd, $J = 13.8$ Hz, $J = 5.3$ Hz, 1H), 3.77-3.22 (m, 1H), 3.06 (t, $J = 7.5$ Hz, 4 H), 2.79 (d, $J = 10.0$ Hz, 1H), 2.66 (d, $J = 9.9$ Hz, 1H), 2.04 (t, $J = 6.5$ Hz, 2H).

Example 4: (1,4-Diazabicyclo[3.2.1]oct-4-yl)-(5-phenyl-furan-2-yl)-methanone



15 By the process described in Example 1, diazabicyclo[3.2.1]octane dihydrochloride salt was reacted with 5-phenyl-furan-2-carboxylic acid to afford the title compound as a white solid. MS (APCI+) 283 $[M+1]^+$. ^1H NMR (300.132 MHz, CDCl_3) δ 7.68 - 7.62 (m, 2H), 7.50 - 7.32 (m, 4H), 6.77 (bs, 1H), 5.56 (m, 1H), 4.72 (m, 1H), 3.72 (m, 2H), 3.38 (m, 5H), 2.61 - 2.43 (m, 1H), 2.38 - 2.20 (m, 1H).


20 **Example 5:** Benzofuran-2-yl-(1,4-diazabicyclo[3.2.1]oct-4-yl)-methanone



By the process described in Example 1, diazabicyclo[3.2.1]octane dihydrochloride salt was reacted with 2-benzofurancarboxylic acid to afford the title compound as an off-white

solid (34 mg, 60 %). MS (APCI+) 257 [M+1]⁺. ¹H NMR (300.132 MHz, CDCl₃) δ 7.68 (d, *J* = 7.7 Hz, 1H), 7.53 (d, *J* = 8.3 Hz, 1H), 7.44 (t, *J* = 6.5 Hz, 1H), 7.42 (s, 1H), 7.32 (t, *J* = 7.5 Hz, 1H), 5.39 (s, 1H), 4.49 (dd, *J* = 4.5 Hz, *J* = 14.4 Hz, 1H), 3.67 (quintet, *J* = 6.7 Hz, 1H), 3.53 (sextet, *J* = 6.0 Hz, 1H), 3.44 - 3.02 (m, 4H), 2.42-2.14 (m, 2H), 1.61 - 1.54 (m, 1H).


Example 6: (1,4-Diazabicyclo[3.2.1]oct-4-yl)-(1-methyl-1*H*-indol-2-yl)-methanone



By the process described in Example 1, diazabicyclo[3.2.1]octane dihydrochloride salt was reacted with 3*H*-indole-2-carboxylic acid to afford the title compound as a colorless gum.

MS (APCI+) 270 [M+1]⁺. ¹H NMR (300.132 MHz, CDCl₃) δ 7.62 (d, *J* = 8.0 Hz, 1H), 7.35 (t, *J* = 8.9 Hz, 1H), 7.28 (d, *J* = 8.0 Hz, 1H), 7.14 (t, *J* = 7.5 Hz, 1H), 6.59 (s, 1H), 5.32-4.67 (m, 1H), 4.31-3.78 (m, 1H), 3.84 (s, 3H), 3.08 (d, *J* = 7.7 Hz, 1H), 3.05 (t, *J* = 7.2 Hz, 3H), 2.88 - 2.71 (m, 1H), 2.69 - 2.54 (m, 1H), 1.99 (m, 2H).


Example 7: Biphenyl-3-yl-(1,4-diazabicyclo[3.2.1]oct-4-yl)-methanone



By the process described in Example 1, diazabicyclo[3.2.1]octane dihydrochloride salt with biphenyl-3-carboxylic acid to afford the title compound as a gum. MS (APCI+) 293

[M+1]⁺. ¹H NMR (300.132 MHz, CDCl₃) δ 7.68 - 7.56 (m, 4H), 7.49 (d, *J* = 8.0 Hz, 1H), 7.45 (t, *J* = 7.4 Hz, 2H), 7.37 (t, *J* = 7.5 Hz, 2H), 5.24 (bs, 1H), 5.24 (bs, 1H), 3.41 (bs, 1H), 3.13 - 2.95 (m, 4H), 2.95 - 2.43 (m, 2H), 2.18 - 1.68 (m, 2H).


Example 8: (1,4-Diazabicyclo[3.2.1]oct-4-yl)-(4-methoxy-phenyl)-methanone



By the process described in Example 1, diazabicyclo[3.2.1]octane dihydrochloride salt was reacted with 4-methoxy-benzoic acid to afford the title compound as an off-white film.


MS (APCI+) 247 [M+1]⁺. ¹H NMR (300.132 MHz, CDCl₃) δ 7.38 (d, *J* = 8.5 Hz, 2H), 6.92 (d, *J* = 8.7 Hz, 2H), 3.84 (s, 3H), 3.29 (m, 1H), 3.07 (d, *J* = 10.2 Hz, 4H), 2.86 - 2.73 (m, 2H), 2.66 (d, *J* = 10.7 Hz, 2H), 2.00 (m, 2H).

Example 9: (1,4-Diazabicyclo[3.2.1]oct-4-yl)-(1*H*-indol-5-yl)-methanone



By the process described in Example 1, diazabicyclo[3.2.1]octane dihydrochloride salt was reacted with 3*H*-indole-5-carboxylic acid to afford the title compound as an off-white film. MS (APCI+) 256 [M+1]⁺. ¹H NMR (300.132 MHz, CDCl₃) δ 8.36 (bs, 1H), 7.72 (s, 1H), 7.41 (d, *J* = 8.4 Hz, 1H), 7.26 (m, 1H), 6.60 (s, 1H), 3.70 - 3.16 (m, 1H), 3.09 (d, *J* = 12.5 Hz, 2H), 3.04 (t, *J* = 8.0 Hz, 2H), 2.85 - 2.66 (m, 1H), 2.66 - 2.52 (m, 1H), 1.98 (m, 2H), 1.70 (m, 2H).


Example 10: (1,4-Diazabicyclo[3.2.1]oct-4-yl)-naphthalen-2-yl-methanone



By the process described in Example 1, diazabicyclo[3.2.1]octane dihydrochloride salt was reacted with naphthalene-2-carboxylic acid to afford the title compound as an amber gum. MS (APCI+) 267 [M+1]⁺. ¹H NMR (300.132 MHz, CDCl₃) δ 7.88 (d, *J* = 8.0 Hz, 2H), 7.87 (t, *J* = 6.5 Hz, 2H), 7.54 (m, 2H), 7.48 (dd, *J* = 8.3, 1.3 Hz, 1H), 5.26 (m, 1H), 4.30 (m, 1H), 3.43 (m, 1H), 3.09 (d, *J* = 12.0 Hz, 2H), 3.05 (m, 2H), 2.88 (m, 1H), 2.75 - 2.46 (m, 1H), 2.00 (s, 2H).

CLAIMS

1. A compound according to Formula I:



- 5 wherein:

D is selected from oxygen, sulfur or $N(R^1)_2$;

- Ar^1 is selected from a 5- or 6-membered aromatic or heteroaromatic ring having 0, 1 or 2 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms, or selected from an 8-, 9- or 10-membered fused aromatic or heteroaromatic ring system having 0, 1, 2 or 3 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms;

E is a single bond, -O-, -S-, or - NR^2 ;

G is selected from hydrogen, C_1 - C_4 alkoxy or Ar^2 , where Ar^2 is a 5- or 6-membered aromatic or heteroaromatic ring having 0, 1 or 2 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms;

- 15 where each Ar^1 or Ar^2 moiety independently is unsubstituted or has 1, 2 or 3 substituents selected from - R^3 , - C_1 - C_6 alkyl, - C_2 - C_6 alkenyl, - C_2 - C_6 alkynyl, halogen, -CN, - NO_2 , - CF_3 , - $S(O)_nR^3$, - NR^2R^3 , - $CH_2NR^2R^3$, - OR^3 , - CH_2OR^3 or - CO_2R^4 ;

R^1 , R^2 and R^3 are independently selected at each occurrence from hydrogen, - C_1 - C_4 alkyl, aryl, heteroaryl, - $C(O)R^4$, - $C(O)NHR^4$, - CO_2R^4 or - SO_2R^4 , or

- 20 R^2 and R^3 in combination is $-(CH_2)_jG(CH_2)_k-$ wherein G is oxygen, sulfur, NR^4 , or a bond;

j is 2, 3 or 4;

k is 0, 1 or 2;

n is 0, 1 or 2, and

- 25 R^4 is independently selected at each occurrence from hydrogen, - C_1 - C_4 alkyl, aryl, or heteroaryl,

and stereoisomers, enantiomers, *in vivo*-hydrolysable precursors or a pharmaceutically-acceptable salt thereof.

2. A compound according to Claim 1, wherein:

D is oxygen;

Ar¹ is selected from phenyl or a 5-membered heteroaromatic ring having 0 or 1 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms, or selected from a 9-membered fused aromatic or heteroaromatic ring system having 0, 1, 2 or 3 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms;;

wherein:

E is a single bond;

G is selected from hydrogen, methoxy or Ar², where Ar² is selected from a 6-membered aromatic or heteroaromatic ring having 0 or 1 nitrogen atoms, 0 or 1 oxygen atoms, and 0 or 1 sulfur atoms;

where each Ar¹ or Ar² moiety independently is unsubstituted or has 1, 2 or 3 substituents selected from halogen, -CN, -NO₂, -CF₃, -CH₃ or -C₂H₅;

and stereoisomers, enantiomers, *in vivo*-hydrolysable precursors or a pharmaceutically-acceptable salt thereof.

3. A compound according to Claim 1, wherein:

D is oxygen;

Ar¹ is selected from phenyl, furanyl, thiophenyl or 1-methyl-1H-pyrrolyl:

E is a single bond;

G is selected from hydrogen, methoxy, phenyl or pyridyl, and

Ar¹ bears 1 halogen substituent;

and stereoisomers, enantiomers, *in vivo*-hydrolysable precursors or a pharmaceutically-acceptable salt thereof.

4. A compound according to Claim 1, wherein:

E represents a single bond; or an enantiomer thereof, or a pharmaceutically-acceptable salt thereof.

5. A compound according to Claim 1, wherein:

Ar¹ is furanyl or thiophenyl having optional substituents as defined herein.

6. A compound according to Claim 1, selected from:
(1,4-diazabicyclo[3.2.1]oct-4-yl)-(5-pyridin-3-yl-thiophen-2-yl)-methanone;
(1,4-diazabicyclo[3.2.1]oct-4-yl)-(5-phenyl-thiophen-2-yl)-methanone;
[5-(4-chloro-phenyl)-furan-2-yl]-(1,4-diazabicyclo[3.2.1]oct-4-yl)-methanone;
- 5 (1,4-diazabicyclo[3.2.1]oct-4-yl)-(5-phenyl-furan-2-yl)-methanone;
benzofuran-2-yl-(1,4-diazabicyclo[3.2.1]oct-4-yl)-methanone;
(1,4-diazabicyclo[3.2.1]oct-4-yl)-(1-methyl-1*H*-indol-2-yl)-methanone;
biphenyl-3-yl-(1,4-diazabicyclo[3.2.1]oct-4-yl)-methanone;
(1,4-diazabicyclo[3.2.1]oct-4-yl)-(4-methoxy-phenyl)-methanone;
- 10 (1,4-diazabicyclo[3.2.1]oct-4-yl)-(1*H*-indol-5-yl)-methanone, or
(1,4-diazabicyclo[3.2.1]oct-4-yl)-naphthalen-2-yl-methanone
or stereoisomers, enantiomers, *in vivo*-hydrolysable precursors or a pharmaceutically-acceptable salt thereof.
- 15 7. A method of treatment or prophylaxis of a disease or condition in which activation of the $\alpha 7$ nicotinic receptor is beneficial which method comprises administering a therapeutically-effective amount of a compound according to Claim 1 to a subject suffering from said disease or condition.
- 20 8. The method of Claim 7, wherein said disease or condition is anxiety, schizophrenia, mania or manic depression.
9. A method of treatment or prophylaxis of neurological disorders, psychotic disorders or intellectual impairment disorders, which comprises administering a therapeutically effective
- 25 amount of a compound according to Claim 1.
10. The method of Claim 9, wherein said disorder is Alzheimer's disease, learning deficit, cognition deficit, attention deficit, memory loss, Attention Deficit Hyperactivity Disorder, Parkinson's disease, Huntington's disease, Tourette's syndrome, neurodegenerative disorders
- 30 in which there is loss of cholinergic synapses, jetlag, nicotine addiction, craving, pain, or ulcerative colitis.

11. A method for inducing the cessation of smoking comprising administering an effective amount of a compound according to Claim 1.
12. A pharmaceutical composition comprising a compound according to Claim 1 and a pharmaceutically-acceptable diluent, lubricant or carrier.
13. A method of treatment or prophylaxis of a disease or condition in which activation of the $\alpha 7$ nicotinic receptor is beneficial which method comprises administering a therapeutically-effective amount of a pharmaceutical composition according to Claim 12 to a subject suffering from said disease or condition.
14. The method of Claim 13, wherein said disease or condition is anxiety, schizophrenia, mania or manic depression.
15. A method of treatment or prophylaxis of neurological disorders, psychotic disorders or intellectual impairment disorders, which comprises administering a therapeutically effective amount of a pharmaceutical composition according to Claim 12.
16. The method of Claim 10, wherein said disorder is Alzheimer's disease, learning deficit, cognition deficit, attention deficit, memory loss, Attention Deficit Hyperactivity Disorder, Parkinson's disease, Huntington's disease, Tourette's syndrome, neurodegenerative disorders in which there is loss of cholinergic synapses, jetlag, nicotine addiction, craving, pain, or ulcerative colitis.
17. A method for inducing the cessation of smoking comprising administering an effective amount of a pharmaceutical composition according to Claim 12.
18. The use of a compound according to Claim 1, an enantiomer thereof or a pharmaceutically-acceptable salt thereof, in the manufacture of a medicament for the treatment or prophylaxis of human diseases or conditions in which activation of the $\alpha 7$ nicotinic receptor is beneficial selected from neurological disorders, psychotic disorders, intellectual impairment disorders, Alzheimer's disease, learning deficit, cognition deficit,

attention deficit, memory loss, Attention Deficit Hyperactivity Disorder, anxiety, schizophrenia, mania or manic depression, Parkinson's disease, Huntington's disease, Tourette's syndrome, or neurodegenerative disorders in which there is loss of cholinergic synapses.

5


19. The use of a compound according to Claim 1, in the manufacture of a medicament for the treatment or prophylaxis of jetlag, pain, or ulcerative colitis or to facilitate the cessation of smoking or the treatment of nicotine addiction or craving including that resulting from exposure to products containing nicotine.

10

ABSTRACT

Title: NICOTINIC ACETYLCHOLINE RECEPTOR LIGANDS

5 Acetylcholine receptor ligands of formula I



wherein D, Ar¹, E and Ar² are as described in the specification, diastereoisomers, enantiomers, pharmaceutically-acceptable salts, methods of making, pharmaceutical compositions containing and methods for using the same.